Author Index

Achs, E., see Ulberth, F. 202
Akporhonor, E. E.
——, Le Vent, S. and Taylor, D. R.
Calculation of programmed temperature gas chromatographic characteristics from isothermal data. III. Predicted retention indices and equivalent temperatures 269
Anacleto, J. F., see D'Agostino, P. A. 259
Apps, P. J.
High-precision sampling of trace gas-borne volatiles by the dynamic solvent effect with a comparative review of alternative techniques 21
Arancibia, E. L., see Castells, R. C. 45
Arrigoni-Martelli, E., see Bruno, G. 319
Barbaro, A. M., see Biagi, G. L. 163
Barra, D., see Simmaco, M. 129
Berger, M., see Wagner, J. R. 191
Betts, T. J.
Implications of solvent selectivity triangles in assessing stationary phases for gas chromatography 186
Biagi, G. L.
——, Barbaro, A. M., Guerra, M. C., Borea, P. A. and Recanatini, M.
Lipophilic character of cardiac glycosides. R_m values as lipophilicity parameters 163
Biase, D. de, see Simmaco, M. 129
Bonner, W. A.
—— and Lee, S. Y.
Quantitative gas chromatographic analysis of mandelic acid enantiomers 287
Borea, P. A., see Biagi, G. L. 163
Bossa, F., see Simmaco, M. 129
Bronzetti, M., see Bruno, G. 319
Brooks, P. W., see D'Agostino, P. A. 259
Bruno, G.
———, Gasparrini, F., Misiti, D., Arrigoni-Martelli, E. and Bronzetti, M.
High-performance liquid chromatographic separation of biomolecules using calcium phosphate supported on macroporous silica microparticles 319
Bruyn, A. De, see Van den Eeckhout, E. 113
Cadet, J., see Wagner, J. R. 191
Cappiello, P. E.
—— and Kling, G. J.
Determination of zeatin and zeatin riboside in plant tissue by solid-phase extraction and ion-exchange chromatography 197
Castells, R. C.
——, Arancibia, E. L. and Nardillo, A. M.
Regression against temperature of gas chromatographic retention data 45
Cauwelaert, F. Van, see Malfait, T. 369
Chen, Y.-Y., see Wu, C.-Y. 279
Churáček, J., see Jandera, P. 297
Ciambra, C. M., see Karp, S. 434
Claereboudt, J., see Van den Eeckhout, E. 113
Claeys, M., see Van den Eeckhout, E. 113
Cooper, D. A., see Lurie, I. S. 391
DaCunha, A. R., see Stout, S. J. 429
D'Agostino, P. A.
——, Provost, L. R., Anacleto, J. F. and Brooks, P. W.
Capillary column gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry detection of chemical warfare agents in a complex airborne matrix 259
Daučik, P., see Drdák, M. 207
De Biase, D., see Simmaco, M. 129
De Bruyn, A., see Van den Eeckhout, E. 113
De Ligny, C. L., see Tji, T. G. 403
Derksen, R. J. A. M., see Gorris, L. G. M. 421
Drdák, M.
——, Daučík, P. and Kubaský, J.
Analysis of anthocyanins in red wines by high-performance liquid chromatography using butylamines in the mobile phase 207
Drift, C. van der, see Gorris, L. G. M. 421
Eeckhout, E. Van den, see Van den Eeckhout, E. 113
El Tayar, N., see Vallat, P. 411
Englert, D., see Tsai, A. M. 89
Esmans, E. L., see Van den Eeckhout, E. 113
Ferrara, P., see Marchese, E. 351
Fischer, N. H., see Weidenhamer, J. D. 151
Frankenberger, Jr., W. T.
——, Mehra, H. C. and Gjerde, D. T.
Environmental applications of ion chromatography (Review) 211
Fréchet, J. M. J., see Rolls, W. A. 97
García-Pérez, A.-I.
——, Sancho, P. and Luque, J.
Settling-time dependence of rat bone marrow cell partition and counter-current distribution in charge-sensitive aqueous two-phase systems. Relationship with the cell partitioning mechanism 79
Gasparrini, F., see Bruno, G. 319
Gelsema, W. J., see Tji, T. G. 403
Giddings, J. C.
Two-dimensional field-flow fractionation 247

Gjerde, D. T., see Frankenberger, Jr., W. T. 211
Gorris, L. G. M.
——, Korteland, J., Derksen, R. J. A. M., Van der Drift, C. and Vogels, G. D.
Quantification of cyclic 2,3-diphosphoglycerate from methanogenic bacteria by isotachophoresis 421
Graham, E. E., see Tsai, A. M. 89
Gray, D. O., see Grosvenor, P. W. 456
Grosvenor, P. W.
—— and Gray, D. O.
2,4-Dinitrophenylpyridium chloride, a novel and versatile reagent for the detection of amino acids, primary and secondary amines, thiols, thiolactones and carboxylic acids during planar chromatography 456
Guerra, M. C., see Biagi, G. L. 163
Guiochon, G., see Huang, J.-X. 335
Hostettmann, K., see Vallat, P. 411
Hruby, V. J., see Toth, G. 450
Huang, J.-X.
——, Schudel, J. and Guiochon, G.
Absorption behaviour of albumin and conalbumin on TSK-DEAE 5 PW anion exchanger 335
Iwaki, K.
——, Okumura, N., Yamazaki, M., Nimura, N. and Kinoshita, T.
Precolumn derivatization technique for high-performance liquid chromatographic determination of penicillins with fluorescence detection 359
Jandera, P.
——, Urbánek, J., Prokeš, B. and Churáček, J.
Comparison of various stationary phases for normal-phase high-performance liquid chromatography of ethoxylated alkylphenols 297
Jones, H. K.
——, Nguyen, N. T. and Smith, R. D.
Variance contributions to band spread in capillary zone electrophoresis 1
Jordan, E. D., see Weidenhamer, J. D. 151
Karp, S.
——, Ciambra, C. M. and Miklean, S.
High-performance liquid chromatographic post-column reaction system for the electrochemical detection of ascorbic acid and dehydroascorbic acid 434
Kaufmann, G. B., see Ray, R. K. 464
Kinoshita, T., see Iwaki, K. 359
Kling, G. J., see Cappiello, P. E. 197
Korteland, J., see Gorris, L. G. M. 421
Kram, T. C., see Lurie, I. S. 391
Krips, H. J., see Tji, T. G. 403
Kubaský, J., see Drdák, M. 207
Lampert, B. M.
—— and Stewart, J. T.
Determination of non-steroidal anti-inflammatory analgesics in solid dosage forms by high-performance liquid chromatography on underivatized silica with aqueous mobile phase 381
Lebl, M., see Toth, G. 450
Lee, S. Y., see Bonner, W. A. 287
Lee, W.-C.
——, Tsai, G. J. and Tsao, G. T.
Effects of isotherm non-linearity on the determination of the binding constant in affinity chromatography 55
Le Vent, S., see Akporhonor, E. E. 269
Li, H.-Y., see Wu, C.-Y. 279
Lier, J. E. van, see Wagner, J. R. 191
Ligny, C. L. de, see Tji, T. G. 403
Lu, X.-R., see Wu, C.-Y. 279
Luque, J., see García-Pérez, A.-I. 79
Lurie, I. S.
——, Moore, J. M., Kram, T. C. and Cooper, D. A.
Isolation, identification and separation of isomeric truxillines in illicit cocaine 391
Malfait, T.
—— and Van Cauwelaert, F.
Preparative and analytical separation of oligosaccharides from κ-carrageenan 369
Marchese, E.
——, Vita, N., Maureaud, T. and Ferrara, P.
Separation by cation-exchange high-performance liquid chromatography of three forms of Chinese hamster ovary cell-derived recombinant human interleukin-2 351
Marston, A., see Vallat, P. 411
Maruyama, T., see Takeba, K. 441
Matsumoto, U.
——, Shibusawa, Y. and Yamashita, M.
Surface affinity chromatographic separation of blood cells. VII. Relationship between capacity factors of human peripheral blood cells and the rate of penetration of liquids into xerogel column packings 69
Matsumoto, M., see Takeba, K. 441
Maureaud, T., see Marchese, E. 351
Mehra, H. C., see Frankenberger, Jr., W. T. 211
Miklean, S., see Karp, S. 434
Misiti, D., see Bruno, G. 319
Moore, J. M., see Lurie, I. S. 391
Nakazawa, H., see Okamoto, M. 445
——, see Takeba, K. 441
Nardillo, A. M., see Castells, R. C. 45
Nguyen, N. T., see Jones, H. K. 1
Nimura, N., see Iwaki, K. 359

Okamoto, M.
—— and Nakazawa, H.
Direct liquid chromatographic resolution of (*R*)- and (*S*)-abscisic acid using a chiral ovomucoid column 445
Okumura, N., see Iwaki, K. 359
Palmqvist, B., see Petersen, T. G. 139
Pepermans, H., see Van den Eeckhout, E. 113
Petersen, T. G.
—— and Palmqvist, B.
Utilizing column selectivity in developing a high-performance liquid chromatographic method for ginsenoside assay 139
Prokeš, B., see Jandera, P. 297
Provost, L. R., see D'Agostino, P. A. 259
Ray, R. K.
—— and Kaufmann, G. B.
Paper chromatographic studies of metal complexes. II. Comparison of square planar and octahedral complexes 464
Recanatini, M., see Biagi, G. L. 163
Rolls, W. A.
—— and Fréchet, J. M. J.
High-performance liquid chromatography separation media based on functional polymers containing phenolic hydroxyls 97
Saitoh, K., see Suzuki, N. 179
Sancho, P., see García-Pérez, A.-I. 79
Schudel, J., see Huang, J.-X. 335
Shibata, Y., see Suzuki, N. 179
Shibusawa, Y., see Matsumoto, U. 69
Simmaco, M.
——, De Biase, D., Barra, D. and Bossa, F.
Automated amino acid analysis using precolumn derivatization with dansylchloride and reversed-phase high-performance liquid chromatography 129
Sinsheimer, J. E., see Van den Eeckhout, E. 133
Slacanin, I., see Vallat, P. 411
Smith, R. D., see Jones, H. K. 1
Stewart, J. T., see Lampert, B. M. 381
Stout, S. J.
—— and DaCunha, A. R.
Simple, direct gas chromatography–mass spectrometry interface for the ion trap detector 429
Suzuki, N.
——, Saitoh, K. and Shibata, Y.
High-performance thin-layer chromatography of rare earth tetraphenylporphine complexes 179
Takeba, K.
——, Maruyama, T., Matsumoto, M. and Nakazawa, H.
Determination of tyramine in cheese by reversed-phase high-performance liquid chromatography with amperometric detection 441
Tayar, N. El, see Vallat, P. 411
Taylor, D. R., see Akporhonor, E. E. 269
Testa, B., see Vallat, P. 411
Tji, T. G.
——, Krips, H. J., Gelsema, W. J. and De Ligny, C. L.
Determination of the charge of ions by partition coefficient measurements in gel permeation chromatography 403
Toth, G.
——, Lebl, M. and Hruby, V. J.
Chiral thin-layer chromatographic separation of phenylalanine and tyrosine derivatives 450
Tsai, A. M.
——, Englert, D. and Graham, E. E.
Study of the dynamic binding capacity of two anion exchangers using bovine serum albumin as a model protein 89
Tsai, G. J., see Lee, W.-C. 55
Tsao, G. T., see Lee, W.-C. 55
Ulberth, F.
—— and Achs, E.
Argentation chromatography of fatty acid methyl esters using silver-loaded solid-phase extraction columns 202
Urbánek, J., see Jandera, P. 297
Vallat, P.
——, El Tayar, N., Testa, B., Slacanin, I., Marston, A. and Hostettmann, K.
Centrifugal counter-current chromatography, a promising means of measuring partition coefficients 411
Van Cauwelaert, F., see Malfait, T. 369
Van den Eeckhout, E.
——, De Bruyn, A., Pepermans, H. Esmans, E. L., Vryens, I., Claereboudt, J., Claeys, M. and Sinsheimer, J. E.
Adduct formation identification between phenyl glycidyl ether and 2′-deoxyadenosine and thymidine by chromatography, mass spectrometry and nuclear magnetic resonance spectroscopy 113
Van der Drift, C., see Gorris, L. G. M. 421
Van Lier, J. E., see Wagner, J. R. 191
Vent, S. le, see Akporhonor, E. E. 269
Vita, N., see Marchese, E. 351
Vogels, G. D., see Gorris, L. G. M. 421
Vryens, I., see Van den Eeckhout, E. 113
Wagner, J. R.
——, Berger, M., Cadet, J. and Van Lier, J. E.
Analysis of thymidine hydroperoxides by post-column reaction high-performance liquid chromatography 191

Weidenhamer, J. D.
——, Jordan, E. D. and Fischer, N. H. Evaluation of high-performance liquid and capillary gas chromatography for analysis of sesquiterpene lactones of the Melampodiinae 151

Wu, C.-Y.
——, Li, H.-Y., Chen. Y.-Y. and Lu, X.-R. Preparation and gas chromatographic characterization of some immobilized crown ether–polysiloxane stationary phases 279

Yamashita, M., see Matsumoto, U. 69
Yamazaki, M., see Iwaki, K. 359

Errata

J. Chromatogr., 405 (1987) 67–76.
Page 70, 3rd line, "eqn.4" should read 'eqn. 4a".
Page 70, 4th line from bottom, "$\exp(k_2 T)$" should read "$\exp(k_2/T)$".
Page 71, last line, "$k_{3(1)}$" should read "$k_{3(1)}t_1$".
Page 73, eqn. 13, "$I_{eq(2)}$" should read "$T_{eq(2)}$".
Page 74, 9th line, "$\bar{t}$" should read "T".

J. Chromatogr., 503 (1990) 155–165.
Page 156, last two text lines, sentence "B_1 and B_2 were stainless-steel pipes, 1.1 m in length and of 0.5 mm I.D." should be deleted.